

Europæisk standardserie - anbefalede minimale testsæt.
(Allergener fra Hermal Trolab eller Chemotechnique)

Allergener	Konc./vehikel	Alternativ
Potassium dichromate	0,5% pet.	
Neomycin sulfate	20% pet.	
Thiuram mix	1% pet.	
p-phenylenediamine	1% pet.	
Cobalt chloride	1% pet.	
Benzocaine	5% pet.	
Formaldehyde	1% aq.	
Colophonium	20% pet.	
Clioquinol	5% pet.	Quinoline mix 6% in pet.
Myroxylon pereirae (<i>Balsam of Peru</i>)	25% pet.	
N-Isopropyl-N'-phenylparaphenylenediamine (<i>IPPD</i>)	0,1% pet.	
Wool alcohols (<i>lanolin alcohol</i>)	30% pet.	
Mercapto mix	1% pet.	
Epoxy resin	1% pet.	
Paraben mix	16% pet.	
para-tertiary-butylphenol-formaldehyde resin (<i>PTBP resin</i>)	1% pet.	
Fragrance mix I	8% pet.	
Quaternium-15	1% pet.	
Nickel sulphate	5% pet.	
Methylisothiazolinone/methylchlorisothiazolinone	0,01% aq.	
Mercaptobenzothiazole	2% pet.	
Primin	0,01% pet.	
Sesquiterpene lactone mix	0,1% pet.	
Budesonide	0,01% pet.	
Tixocortol pivalate	0,1% pet.	
Hydroxyisohexyl-3-cyclohexene carboxaldehyde (<i>Lyrall</i>)	5% pet.	
Methyldibromo glutaronitrile	0,5% pet.	
Fragrance mix II	14 % pet.	
<i>Kilde: Andersen KE. Textbook Contact Dermatitis 4 th. Kap. 29; Bruze M. Contact Dermatitis 2005:52:24-8; Frosch PJ. Contact Dermatitis 2005:52.4:207-15</i>		

Anvendes **True Test** panel 1 og 2 bør testen, som et minimum suppleres med:

Primin	0,01% pet.
Sesquiterpene lactone mix	0,1% pet.
Budesonide	0,01% pet.
Tixocortol pivalate	0,1% pet.
Hydroxyisohexyl-3-cyclohexene carboxaldehyde (<i>Lyrall</i>)	5% pet.
Methyldibromo glutaronitrile	0,5% pet.
Fragrance mix II	14 % pet.

I TRUE Testen indgår black rubber mix i stedet for IPPD og caine mix i stedet for benzocaine.

Juli 2007