

Europæisk basisserie - anbefalede minimale testsæt.
(Allergener i petrolatum/vand)

Allergener	Konc./vehikel	Alternativ
Potassium dichromate	0,5% pet.	
Neomycin sulfate	20% pet.	
Thiuram mix	1% pet.	
p-phenylenediamine	1% pet.	
Cobalt chloride	1% pet.	
Benzocaine	5% pet.	
Formaldehyde	2% aq.	
Colophonium	20% pet.	
Clioquinol	5% pet.	Quinoline mix 6% in pet.
Myroxylon pereirae (<i>Balsam of Peru</i>)	25% pet.	
N-Isopropyl-N'-phenylparaphenylenediamine (<i>IPPD</i>)	0,1% pet.	
Wool alcohols (<i>lanolin alcohol</i>)	30% pet.	
Mercapto mix	1% pet.	
Epoxy resin	1% pet.	
Paraben mix	16% pet.	
para-tertiary-butylphenol-formaldehyde resin (<i>PTBP resin</i>)	1% pet.	
Fragrance mix I	8% pet.	
Quaternium-15	1% pet.	
Nickel sulphate	5% pet.	
Methylisothiazolinone/methylchlorisothiazolinone	0.02% aq.	
Mercaptobenzothiazole	2% pet.	
Primin	0,01% pet.	
Sesquiterpene lactone mix	0,1% pet.	
Budesonide	0,01% pet.	
Tixocortol pivalate	0,1% pet.	
Hydroxyisohexyl-3-cyclohexene carboxaldehyde (<i>Lyrall</i>)	5% pet.	
Methyldibromo glutaronitrile	0,5% pet.	
Fragrance mix II	14 % pet.	
Methylisothiazolinone	0,2% aq.	
Textile mix	6,6% pet	

Kilde: Bruze M, Goossens A, Isaksson M. Contact Dermatitis 2014;71, 35–40; Bruze M et al 2015: in press

Anvendes **True Test panel 1, 2 og 3** bør testen, som et minimum suppleres med:

Sesquiterpene lactone mix	0,1% pet.
Hydroxyisohexyl-3-cyclohexene carboxaldehyde (<i>Lyrall</i>)	5% pet.
Fragrance mix II	14 % pet.
Methyldibromo glutaronitrile	0,5% pet.
Methylisothiazolinone	0,2% aq.
Methylisothiazolinone/methylchlorisothiazolinone	0,02% aq
Textile mix	6,6% pet.
Primin	0,01% pet.
Formaldehyde	2% aq.

I TRUE Testen indgår black rubber mix i stedet for IPPD og caine mix i stedet for benzocaine.

Marts 2015